

References

- [1] ISIS-2 (SECOND INTERNATIONAL STUDY OF INFARCT SURVIVAL) COLLABORATIVE GROUP (1988) 'Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17 187 cases of suspected acute myocardial infarction: isis-2', *The Lancet*, 2(8607), pp. 349-360.
- [2] Bolooki, H. and Askari, A. (2010) *Acute Myocardial Infarction*. Available at: <http://www.clevelandclinicmeded.com/medicalpubs/diseasemanagement/cardiology/acute-myocardial-infarction/> (Accessed 13 November 2016).
- [3] Brookes, S., Whitley, E., Peters, T., Mulheran, P., Egger, M. and Davey Smith, G. (2001) 'Subgroup analyses in randomised controlled trials: quantifying the risks of false-positives and false-negatives', *Health Technology Assessment*, 5(33), pp. 1-56.
- [4] The Canadian Cooperative Study Group (1978) 'A Randomized Trial of Aspirin and Sulfinpyrazone in Threatened Stroke', *New England Journal of Medicine*, 299(2), pp. 53-59.
- [5] Rothwell, P. (2005) 'Subgroup analysis in randomised controlled trials: importance, indications, and interpretation', *Lancet*, 365(9454), pp. 176–86.
- [6] Rothwell, P., Eliasziw, M., Gutnikov, S., Warlow, C. and Barnett, H. (2004) 'Endarterectomy for symptomatic carotid stenosis in relation to clinical subgroups and timing of surgery', 363(9413), pp. 915-24.
- [7] Kasenda, B., Schandelmaier, S., Sun, X., von Elm, E., You, J., Blümle, A., et al. (2014) 'Subgroup analyses in randomised controlled trials: cohort study on trial protocols and journal publications', *BMJ*, 349, pp. g4539.