

References

1. nih.gov. (2019). NIMH » Anxiety Disorders. [online] Available at: <https://www.nimh.nih.gov/health/topics/anxiety-disorders/index.shtml> [Accessed 12 May 2019].
2. Breilmann, J., Girlanda, F., Guaiana, G., Barbui, C., Cipriani, A., Castellazzi, M., Bighelli, I., Davies, S., Furukawa, T. and Koesters, M. (2019). [Benzodiazepines versus placebo for panic disorder in adults](#). *Cochrane Database of Systematic Reviews* 2019, Issue 3. Art No: CD010677. DOI: 10.1002/14651858.CD010677.pub2
3. Pompoli, A., Furukawa, T., Imai, H., Tajika, A., Efthimiou, O. and Salanti, G. (2016). [Psychological therapies for panic disorder with or without agoraphobia in adults: a network meta-analysis](#). *Cochrane Database of Systematic Reviews*. 2016, Issue 4, Art. No: CD011004. DOI: 10.1002/14651858. CD011004.pub2
4. Nice.org.uk. (2019). Guidance | Generalised anxiety disorder and panic disorder in adults: management | Guidance | NICE. CG113. [online] Available at: <https://www.nice.org.uk/guidance/cg113/chapter/1-Guidance#stepped-care-for-people-with-panic-disorder> [Accessed 12 May 2019].
5. Gomez, A., Barthel, A. and Hofmann, S. (2018). [Comparing the efficacy of benzodiazepines and serotonergic anti-depressants for adults with generalized anxiety disorder: a meta-analytic review](#). *Expert Opinion on Pharmacotherapy*, 19(8), pp.883-894. doi: 10.1080/14656566.2018.1472767.